

Dpnl

5'-GATC-3'
3'-CTAG-5'

Cat. No.	size
E2135-01	500 units
E2135-02	2500 units

Reaction Temperature: 37°C

Inactivation Temperature (20 min): 80°C

Prototype: Dpnl

Source: *Diplococcus pneumoniae*

Purified from *E.coli* strain that carries the Dpnl gene from *Diplococcus pneumoniae*.

Package Contents:

- Dpnl
- 10 x ONE Buffer.

Storage Conditions: Store at -20°C.

Double Digestion – Buffer Compatibility:

ONE Buffer is compatible with most EURx restriction enzymes.

The relative activity of Dpnl in Pfu (Cat.No. E1114) and PfuPlus! (Cat.No. E1118) DNA Polymerase buffer is approx. 50%.

Note 1: Dpnl cleaves only methylated GATC sites.

Note 2: It may be necessary to add more enzyme to obtain complete digestion when using other buffer than optimal (ONE Buffer).

Note 3: It is recommended to add 10 U of the enzyme to obtain complete digestion of the methylated template after standard site-directed mutagenesis (total reaction volume is 50 µl).

Double digestions: 100 µg/ml BSA neither inhibits nor promotes Dpnl cleavage.

DNA Methylation:

No Inhibition: All GATC sites with N⁶-Methyladenine.

Standard Reaction Protocol (for 50 µl volume):

Mix the following reaction components:

- 1-2 µg pure DNA or 10 µl PCR product (=~0.1-2 µg DNA)
- 5 µl 10 x ONE Buffer
- 1-2 U Dpnl (use 1 U per µg DNA, < 10% React. Volume!)

Tips: Add enzyme as last component. Mix components well before adding enzyme. After enzyme addition, mix gently by pipetting. Do not vortex. High (excess) amounts of enzyme can greatly speed up the reaction.

add sterile H₂O to 50 µl final volume

Incubate for 1 h at 37°C

To obtain complete digestion of high molecular weight DNA, (e.g. plant genomic DNA), add excess amounts of enzyme and prolong the incubation time.

Stop reaction by alternatively

- Addition of 2.1 µl EDTA pH 8.0 [0.5 M], final 20 mM or
- Heat Inactivation
20 min at 80°C or
- Spin Column DNA Purification
(e.g. EURx PCR/DNA Clean-Up Kit, Cat. No. E3520) or
- Gel Electrophoresis and Single Band Excision
(e.g. EURx Agarose-Out DNA Kit, Cat.No. E3540) or
- Phenol-Chloroform Extraction or Ethanol Precipitation.

Non-optimal buffer conditions:

To compensate for the lack of enzyme activity, increase the amount of enzyme and/ or reaction time accordingly. The following values may serve as orientation:

- Enzyme amount:** Instead of 1 U enzyme, use ~4 U of enzyme in buffers providing 25% rel. activity, ~2 U in 50%, ~1.5 U in 75% or ~1 U in 100%, respectively.
- Reaction time:** Increase by ~1.3-fold (75% rel. activity), ~2-fold (50%) or ~4-fold (25%).

Unit Definition:

One unit is the amount of enzyme required to completely digest 1 µg of pBR322 *dam* methylated DNA in 1 hr. Total reaction volume is 50 µl. Enzyme activity was determined in the recommended reaction buffer.

Reaction Buffer:

1 x ONE Buffer

Storage Buffer:

10 mM Tris-HCl (pH 7.4 at 4°C), 1 mM dithiothreitol, 400 mM NaCl, 0.1 mM EDTA, 0.1% (v/v) Triton™X-100, 200 µg/ml bovine serum albumin and 50%(v/v) glycerol.

Quality Control:

All preparations are assayed for contaminating endonuclease, 3'-exonuclease, as well as nonspecific single- and double-stranded DNase activities.