



## Fast Polar-BAP

*Thermosensitive  
Bacterial Alkaline Phosphatase*

Cat. No.	size
E1028-01	1000 u
E1028-02	5000 u

### Unit Definition:

One unit is the amount of enzyme required to dephosphorylate 1 µg of linearized pUC19 vector DNA in 10 min at 37°C.

### Storage Conditions:

Store at -20°C.

### Inactivation Temperature (5 min): 70°C

### Quality Control:

All preparations are assayed for contaminating endonuclease and nonspecific RNase and single- and double-stranded DNase activities.

### References:

1. Sambrook (1989) *Molecular Cloning: A Laboratory Manual*, (2nd ed.). 5.72.
2. Werle et al. (1994) *Nuc. Acids Research* 22(20): 4354-4355.

Thermosensitive Bacterial Alkaline Phosphatase catalyzes the release of 5'- and 3'-phosphate groups from DNA, RNA, and nucleotides.

### Description:

- Removes 5'-phosphates from DNA, RNA, rNTPs and dNTPs.
- Heat inactivation in 5 minutes at 70°C.
- Resistant to chemical changes and active over a broad range of buffer conditions.
- Degradation of dNTPs prior to sequencing of PCR product.
- Can be used to remove 5'-phosphates from DNA or RNA prior to 5'-end labeling (1).
- Removes 5'-phosphates from linearized vector molecules to prevent self-ligation of the vector during cloning procedures (1).
- Protein dephosphorylation.
- Ideal for diagnostic immunoassays and immunodetection of proteins and nucleic acids following blotting experiments (1).

### Standard PCR Clean-up Protocol:

#### Mix the following reaction components:

- 25-50 µl of PCR just after amplification
- 0.5 µl of Exonuclease I 20 U/µl (Cat. No. E1150)
- 1 µl of Fast Polar-BAP 1 U/µl

**Incubate** for 15 min at 37°C.

**Heat Inactivate** 15 min 80°C.

- Up to 5 µl may be used directly to sequencing without any other purification.
- It is recommended to use PCR without unspecific products.
- No specific buffer required.

### Standard Vector Dephosphorylation Protocol:

#### Mix the following reaction components:

- 1 µg of DNA cut with any restriction enzyme
- 5 µl 10 x Polar-BAP Reaction Buffer
- 1 µl of Fast Polar-BAP 1 U/µl
- Nuclease-free water: to 50 µl

**Incubate** for 10 min at 37°C.

**Heat Inactivate** 5 min 70°C.

### 1 x Polar-BAP Reaction Buffer:

50 mM Bis-Tris-Propane-HCl (pH 6.0 at 25°C), 1 mM MgCl<sub>2</sub>, 0.1 mM ZnCl<sub>2</sub>.

### Storage Buffer:

10 mM Tris-HCl (pH 7.6 at 22°C), 1 mM MgCl<sub>2</sub>, 0.01 mM ZnCl<sub>2</sub>, 1 mM dithiothreitol, 50% glycerol.

This product is developed, designed and sold exclusively for research purposes and in vitro use only.

EURx Ltd. 80-297 Gdańsk Poland ul. Przyrodników 3, NIP 957-07-05-191, KRS 0000202039  
www.eurx.com.pl, orders@eurx.com.pl, tel. +48 58 524 06 97, fax +48 58 341 74 23