



smART RT-qPCR Kit, plus ROX Solution

Cat. No.	size
E0807-01	25 reactions
E0807-02	100 reactions

smART Unit Definition:

One unit incorporates 1 nmol of TTP into acid precipitable material in 10 min at 37°C using poly(A): oligo dT as a template: primer (1).

Storage Conditions: Store at -20°C.

Concentration: 200 U/μl

Quality Control:

All preparations are assayed for contaminating endonucleases, exonucleases, nonspecific RNases, single- and double-stranded DNase activities.

References:

- Houts, G.E., Masakau, M., Ellis, C., Beard, D. and Beard, J.W. (1979) *J. Virol.* 29, 517-522.

smART RT-qPCR Kit, plus ROX Solution is a complete system for efficient synthesis of first strand cDNA and qPCR with ROX dye as passive reference. Kit contains smART Reverse Transcriptase with increased thermostability and processivity. The real time qPCR is performed with easy to use SG qPCR Master Mix (2x).

Description:

- Kit based on processive smART Reverse Transcriptase.
- Increased thermal stability of reverse transcription in the range of 37°C to 65°C.
- Downstream application: qPCR with SYBR Green I and ROX solution in optimized Master Mix (2x) with dATP, dCTP, dGTP, dTTP/dUTP.
- Contains highly efficient Hot Start DNA Polymerase ideal for real time diagnostic.
- Kit contains UNG (uracil-N-glycosylase) to prevent cross contamination.

COMPONENT:	E0807-01	E0807-02
smART (200 U/μl)	25 μl	100 μl
5 x cDNA Buffer	100 μl	400 μl
0.1 M DTT	50 μl	200 μl
Oligo (dT) ₂₀ (50 μM)	25 μl	100 μl
Random hexamers (200 ng/μl)	25 μl	100 μl
10 mM dNTPs Mix	25 μl	100 μl
RNase Inhibitor (50 U/μl)	12.5 μl	50 μl
SG qPCR Master Mix (2x)	350 μl	1.4 ml
ROX Solution, 25 μM	15 μl	60 μl
UNG (uracil-N-glycosylase) (1 U/μl)	6.25 μl	25 μl
RNase-free Water	1 ml	4 x 1 ml

This product is developed, designed and sold exclusively for research purposes and in vitro use only.

EURx Ltd. 80-297 Gdańsk Poland ul. Przyrodników 3, NIP 957-07-05-191, KRS 0000202039
www.eurx.com.pl, orders@eurx.com.pl, tel. +48 58 524 06 97, fax +48 58 341 74 23

I First strand cDNA synthesis:

1. Place 5 x cDNA Buffer at room temperature, thaw and vortex gently.
2. Assamby reaction in RNase-free tube as follows:

Component:	Amount:
RNA (10 ng-5 µg)	x µl
primer*	1 µl
10 mM dNTP Mix	1 µl
RNase-free Water	to 12.5 µl

*50 µM Oligo(dT)₂₀, 200 ng/µl random hexamer primer or 10 µM reverse gene specific primer.

3. *Optional.* If GC-rich or structured RNA template is used, mix gently, centrifuge briefly and incubate 5 min at 65°C then chill on ice.
4. Add the following components to the reaction tube in the indicated order:

Component:	Amount:
5 x cDNA Buffer	4 µl
DTT 0.1 M	2 µl
RNase Inhibitor 50 U/µl	0.5 µl
smART (200 U/µl)	1 µl
Total volume	20 µl

II qPCR with SG qPCR Master Mix (2x) plus ROX Solution

Store SG qPCR Master Mix (2x) at 4°C in the dark. The performance of the master mix is guaranteed up to 9 months. Freezing and thawing of the master mix should be avoided.

- SG qPCR Master Mix (2x) is a universal solution for quantitative real-time PCR and two-step real-time RT-PCR and can be used on most real-time PCR cyclers available.
- The master mix contains Perpetual Taq DNA Polymerase, optimized reaction buffer, dNTPs (dTTP is partially replaced with dUTP) and SYBR Green I dye.
- SYBR Green I enables DNA analysis without having to synthesize expensive sequence-specific labeled probes.
- Perpetual Taq DNA Polymerase contains recombinant Taq DNA Polymerase bound to anti-Taq monoclonal antibodies that block polymerase activity at moderate temperatures.
- The polymerase activity is restored during the initial denaturation step when amplification reactions are heated at 95°C for at least two minutes.
- Use of the "hot start" enzyme prevents extension of misprimed products and primer-dimers during reaction setup leading to higher specificity and sensitivity of PCR reactions.
- The polymerase enables convenient room temperature reaction setup.
- SYBR Green I is a fluorescent dye which binds all double-stranded DNA molecules and emits a fluorescent signal on binding. The excitation and emission maxima of SYBR Green I are at 494 nm and 521 nm, respectively, which are compatible with use on any real-time cycler.
- SG qPCR Master Mix (2x) contains dUTP, which partially replaces dTTP. It allows the optional use of a uracil-N-glycosylase (UNG) to prevent carryover contamination between reactions. UNG removes uracil from any dU-containing contaminating amplicons, leaving abasic sites and making DNA molecules susceptible to hydrolysis during the initial denaturation step.
- The ROX Solution is provided separately. The use of ROX passive reference dye is necessary for all real-time PCR cyclers from Applied Biosystems and optional for cyclers from Stratagene. ROX compensates for variations of fluorescent signal between wells due to slight differences in reaction volume and fluorescence fluctuations. ROX is not involved in PCR reaction, has a different emission spectrum than SYBR Green I and does not interfere with real-time PCR on any instrument. Refer to the table below to determine the recommended amount of ROX (25 µM) required for a specific PCR cycler.

5. Transfer the sample to preheated to appropriate temperature thermal cycler. Incubate as follows:

Oligo(dT) ₂₀ primed:	30-60 min at 50°C (or 37-65°C)
Gene specific primed:	30-60 min at 50°C (or 37-65°C)
Random hexamer primed:	25°C for 10 min, followed by 20-50 min at 50°C (or 37-65°C).

NOTE

50°C is suitable temperature for most targets. For G-C rich RNA templates or with complex secondary structure temperature can be increased to 65°C.

6. Terminate the reaction by incubating at 85°C for 5 min.
7. cDNA is ready for PCR, can be used immediately or stored at -20°C. Use 1-2 µl for 25 µl PCR or qPCR reaction.

This product is developed, designed and sold exclusively for research purposes and in vitro use only.

EURx Ltd. 80-297 Gdańsk Poland ul. Przyrodników 3, NIP 957-07-05-191, KRS 0000202039
www.eurx.com.pl, orders@eurx.com.pl, tel. +48 58 524 06 97, fax +48 58 341 74 23

Recommended amounts of ROX for a specific real-time PCR cyclers

Instrument	Amount of ROX per 25 µl reaction	Amount of ROX per 1.25 ml of 2x master mix	Final ROX concentration
Applied Biosystems: 7300, 7900HT, StepOne, StepOnePlus, ABI PRISM 7000 and 7700	0.5 µl	50 µl	500 nM
Applied Biosystems: 7500, ViiA 7, Stratagene: Mx3000P, Mx3005P, Mx4000	0.5 µl 10 x diluted (in water)	50 µl 10 x diluted (in water)	50 nM
PCR machines from other manufacturers: Bio-Rad, Roche, Corbett, Eppendorf, Cepheid, etc.	Not required	Not required	-

qPCR with SG qPCR Master Mix(2x) plus ROX Solution

- Mix as follows all reagents in 0.2-0.5 ml tube.

Component	Volume/reaction	Final concentration
SG qPCR Master Mix (2x)	12.5 µl	1 x 2.5 mM MgCl ₂
Forward Primer	Variable	0.3-0.5 µM
Reverse Primer	Variable	0.3-0.5 µM
Template cDNA	Variable	1-2 µl
ROX Solution, 25 µM	0.5 µl or 0.5 µl 10 x diluted	500 nM 50 nM
Optional: UNG (uracil-N-glycosylase) 1 U/µl	0.25 µl	0.25 U/reaction
Water, nuclease free	To 25 µl	-
Total volume	25 µl	-

Notes:

- Minimize exposure of SG qPCR Master Mix (2x) and ROX solution to light during handling to avoid loss of fluorescent signal intensity.
- A reaction volume of 25 µl should be used with most real-time cyclers. Other reaction volumes may be used if recommended for a specific instrument.
- Optimal amplicon length in real-time PCR using SYBR Green I is 70-200 bp.
- Thaw if needed, gently vortex and briefly centrifuge all solutions.
- Set up PCR reactions at room temperature. Use of SG qPCR Master Mix (2x) allows room temperature reaction setup.
- Prepare a reaction master mix by adding all the reaction components except template cDNA.
- Mix the reaction mix thoroughly and dispense appropriate volumes into PCR tubes or plates.
- Add template cDNA to the individual PCR tubes or wells containing the reaction mix. For two-step RT-PCR, the volume of cDNA added should not exceed 10% of the final PCR volume.
- Centrifuge briefly to settle down the reaction components and remove bubbles. Bubbles interfere with fluorescent detection.
- Place the samples in the cycler and start the program.
- Standard concentration of MgCl₂ in real-time PCR reaction is 2.5 mM (as provided with the 1 x SG qPCR Master Mix). In most cases this concentration will produce optimal results. However, if a higher MgCl₂ concentration is required, prepare a 25 mM MgCl₂ stock solution and add to a reaction.
- A final primer concentration of 0.3-0.5 µM is usually optimal, but can be individually optimized in range of 0.1 µM to 1 µM. The recommended starting concentration is 0.5 µM. Raising primer concentration may increase PCR efficiency, but negatively affect PCR specificity. Optimal primer concentration depends on the individual reaction and the real-time PCR cycler used.
- Readjust the threshold value for analysis of every run.
- If using Bio-Rad iCycler iQ or MyiQ instruments collect well factors at the beginning of each experiment. Use an external well factor plate according to the manufacturer's recommendations. Factors are used to compensate for any excitation or pipetting variations.

Thermal Cycling Conditions:

Step	Temperature	Time	Number of Cycles
Optional: UNG pre-treatment	50°C	2 min	1
Initial Denaturation	95°C	15 min	1
Denaturation	94°C	15 s	35-45
Annealing	50-60°C	30 s	
Extension	72°C	30 s	
Optional: Data acquisition	X°C	15 s	
Cooling	4°C	Indefinite	1

Notes:

1. The incubation step of 50°C for 2 minutes must be added if a uracil-N-glycosylase is used to prevent carryover contamination. UNG degrades any dUMP-containing PCR products.
2. During the initial denaturation step UNG and antibodies that block Taq DNA Polymerase are inactivated. The anti-Taq antibodies and UNG require at least 2 min or 10 min incubation at 95°C, respectively. When UNG is not used in PCR reaction the duration of the initial denaturation step can be reduced to 2-5 min at 95°C.
3. UNG activity may be partially restored at temperatures lower than 55°C due to refolding. It is recommended to perform PCR using a temperature equal 55°C or above for the annealing step. After completing the PCR cool reactions to 4°C and load directly on a gel or store frozen.
4. Melting curve analysis should be performed to verify the specificity and identity of PCR products. Melting curve analysis is an analysis step built into the software of real-time cyclers. Melting curve data between 65°C and 95°C should be acquired.
5. Data acquisition should be performed during the extension step. To suppress fluorescence readings caused by the generation of primer-dimers an additional data acquisition step can be added to the protocol. It is possible when T_m of primer-dimers differs from T_m of the specific product (T_m are generated during melting curve analysis). The temperature of the data acquisition step should be above T_m of primer-dimers but approximately 3°C below the T_m of the specific product.
6. Always check the PCR product specificity by gel electrophoresis when designing a new assay. Melting temperatures of the specific product and primer-dimers may overlap.

This product is developed, designed and sold exclusively for research purposes and in vitro use only.

EURx Ltd. 80-297 Gdańsk Poland ul. Przyrodników 3, NIP 957-07-05-191, KRS 0000202039
www.eurx.com.pl, orders@eurx.com.pl, tel. +48 58 524 06 97, fax +48 58 341 74 23