

HOT DISCO TAQ DNA POLYMERASE

Concentration: 5 units/ μ l

MADE IN DENMARK

Cat. No.: 257662

2500 Units

-	Hot Disco Taq DNA Polymerase 5 U/ μ l
ID No.	CL0.100-0022
Cap colour	Red
A220006	5 x 100 μ l

Key Features

Hot Disco Taq DNA Polymerase is a modified form of ClearLine Taq DNA Polymerase. A chemical moiety is attached to the enzyme at the active site, which renders the enzyme inactive at room temperature. Thus, during setup and the first ramp of thermal cycling, the enzyme is not active and misprimed primers are not extended. Once the reaction reaches optimal activating temperature, the chemical moiety is cleaved during a 10-15 minutes heat activation step, releasing the active Hot Disco Taq DNA Polymerase into the reaction.

Kit Components

Hot Disco Taq DNA Polymerase in Storage Buffer

5 U/ μ l Hot Disco Taq DNA Polymerase, 20 mM Tris-HCl pH 8.9, 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.5% Tween® 20, 50% glycerol.

Recommended Storage and Stability

Long term storage at -20 °C. Product expiry at -20 °C is stated on the label.

Option: Store at +4 °C for up to 6 months.

Quality Control

Hot Disco Taq DNA Polymerase is tested for contaminating activities, with no traces of endonuclease activity, nicking activity or exonuclease activity.

Unit Definition

One unit is defined as the amount of polymerase that incorporates 10 nmol of dNTPs into acid-precipitable DNA in 30 minutes at 72 °C under standard assay conditions.

Protocol

This protocol serves as a guideline to ensure optimal PCR results when using Hot Disco Taq DNA Polymerase. Optimal reaction conditions such as incubation times, temperatures and amount of template DNA may vary and must be determined individually.

1. Thaw 10x Buffer, dNTP mix and primer solutions. **It is important to thaw the solutions completely (some buffers need to reach room temperature) and mix thoroughly before use to avoid localized concentrations of salts.** Keep all components on ice. The polymerase is provided in glycerol and does not need thawing.

2. Set up reaction mixtures in an area separate from that used for DNA preparation or product analysis. Work on ice at all times.
3. Prepare a master mix according to Table 1. The master mix typically contains all the components needed for extension except the template DNA.

Table 1. Reaction mix and template DNA

Component	Vol./reaction*	Final concentration*
10x Buffer	5 μ l	1x
25 mM MgCl ₂	-	Optional
dNTP mix (10 mM each)	1 μ l	0.2 mM of each dNTP
Primer A (10 μ M)	1 μ l (0.5 – 5 μ l)	0.2 μ M (0.1 – 1.0 μ M)
Primer B (10 μ M)	1 μ l (0.5 – 5 μ l)	0.2 μ M (0.1 – 1.0 μ M)
DNA Pol.	0.4 μ l (0.2 – 1 μ l)	2 units (1 – 5 units)
PCR-grade H ₂ O	X μ l	-
Template DNA	X μ l	genomic DNA: 20 ng (1 – 200 ng) plasmid DNA: 0.5 ng (0.1 – 1 ng) bacterial DNA: 5 ng (1 – 10 ng)
TOTAL volume	50 μl	-

* Suggested starting conditions; theoretically used conditions in brackets. The final volume can be reduced to 25 μ l by using half of the volumes suggested in Vol./reaction, eg. 0.2 μ l Hot Disco Taq instead of 0.4 μ l Hot Disco Taq.

4. Mix the reaction mix thoroughly and dispense appropriate volumes into reaction tubes.
5. Add template DNA to the individual tubes containing the reaction mix.
6. Program the thermal cycler according to the manufacturer's instructions. **Each program must start with an initial heat activation step at 95 °C for 15 minutes.** For maximum yield and specificity, temperatures and cycling times should be optimized for each new template or primer pair.
7. Place the tubes in the thermal cycler and start the reaction.

Table 2. Three-step PCR program

Cycles	Duration of cycle	Temperature
1	15 minutes ^a	95 °C
25 – 35	20 – 30 seconds ^b 20 – 40 seconds ^c 30 – 90 seconds ^d	95 °C 50 – 65 °C 72 °C
1	5 minutes ^e	72 °C

^a. For activation of the hot start enzyme.

^b. Denaturation step: This step is the first regular cycling event and consists of heating the reaction to 95 °C for 20 – 30 seconds. It causes melting of the DNA template by disrupting the hydrogen bonds between complementary bases, yielding single-stranded DNA molecules.

^c. Annealing step: The reaction temperature is lowered to 50 – 65 °C for 20 – 40 seconds allowing annealing of the primers to the single-stranded DNA template. Typically, the annealing temperature is about 3 – 5 °C below the T_m (melting temperature) of the primers used.

^d. Extension/elongation step: Hot Disco Taq polymerase has its optimum activity temperature at 72 °C. At this step the DNA polymerase synthesizes a new DNA strand complementary to the DNA template strand. The extension time depends on the length of the DNA fragment to be amplified. As a rule of thumb, at its optimum temperature the DNA polymerase will polymerize a thousand bases per minute.

^e. Final elongation: This single step is occasionally performed at a temperature of 72 °C for 5 minutes after the last PCR cycle to ensure that any remaining single-stranded DNA is fully extended.

Tween® 20 is a registered trademark of ICI Americas Inc.

Notes:

- For longer DNA targets more DNA polymerase could be added to the PCR master mix.

For Research Use Only. Not for use in diagnostics procedures.

Other product sizes, combinations and customized solutions are available. Please look at www.dutscher.com or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.

Made in Denmark

Issued 08/2021

DOMINIQUE DUTSCHER SAS