

## Catalog Number: HZ-1038

## Data Sheet



Animal Component-Free Human cell expressed Tag-Free

Endot

Endotoxin Free

SC

## Product Description


Animal-free Recombinant Human VEGF 165is expressed in human 293 cells as a homodimeric glycoprotein with an apparent molecular mass of 45 kDa. The E. coli expressed protein is a monomer-dimer mixture with a molecular mass 18 to 34 kDa. The cytokine is produced in a human cell expression system with serum free, chemically defined media. VEGF, also known as vascular permeability factor (VPF) or vasculotropin, is a homodimeric 34-45 kDa, heparin-binding glycoprotein with potent angiogenic, mitogenic, and vascular permeability enhancing activities specific for endothelial cells.

 Alternative Names
 L VEGFA, MVCD1, Vascular permeability factor, VEGF, VEGFA, VEGFA, VPF

 Source
 Human Embryonic Kidney cells (HEK293). HEK293-derived VEGF165 protein

 Species Reactivity
 human,pig

Specifications				
Test	Method	Specification		
Activity	Dose-dependent stimulation of the proliferation of HUVEC cells (Human Umbilical Vein Endothelial Cells)	0.3-3.75 ng/mL		
Molecular Mass	SDS-PAGE	22 to 30 and 50 kDa reduced, 40 to 50 and 75 kDa non-reduced, homodimer, glycosylated		
Purity	SDS-PAGE	>95%		
Endotoxin	LAL	<1 EU/µg		



www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc. 5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

Preparation			
Shipping Temperature	ambient temperature		
Formulation	1x PBS, See Certificate of Analysis for details		
Reconstitution	Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.2 mg/mL in sterile 1x PBS pH 7.4 containing 0.1% endotoxin-free recombinant serum albumin (HSA). Gently swirl or tap vial to mix.		

	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
Stability and Storage	Lyophilized	Room Temperature	2 weeks
, ,	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
		Avoid repeated freeze-thaw cycles.	