




Restriction Enzyme

Apa I




Cat.#	Size	Conc.
FG-Apal	5,000 units	50 units/μl

Store at -20°C

Supplied with: 10X FastGene® Buffer IV (FG-REB4)
10X FastGene® FastCut Buffer (FG-REBHF)
6X DNA Loading Buffer
Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Dilution buffer

FastGene® Diluent A

Heat Inactivation

Apa I can be inactivated at 65°C for 20 min.

Methylation sensitivity

dam methylation: Not sensitive

dcm methylation: Conditionally sensitive

CpG methylation: Conditionally sensitive

Prolonged incubation

A minimum amount of enzyme required to digest 1 μg substrate DNA for 16 hr; 0.13 U.

Relative activity in FastGene® Buffers

FastGene® Buffer I:	100%
FastGene® Buffer II:	25%
FastGene® Buffer III:	0%
FastGene® Buffer IV:	100%
FastGene® FastCut Buffer:	100%

Note

Incubation at 37°C results in 100% activity. However, it is unstable at 37°C with a half-life of only 30 min. Either *dcm* or CpG methylation overlapping its recognition sequence blocks cleavage of DNA. Salt over 50 mM is inhibitory.

Source: *Acetobacter pasteurianus* sub *pasteurianus*

Reaction conditions

1X FastGene® Buffer IV, 25°C
1X FastGene® FastCut Buffer, 25°C

FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 min with FastGene® FastCut Buffer.

1X FastGene® Buffer IV

20 mM Tris-acetate (pH 7.9 at 25°C)
50 mM potassium acetate
10 mM magnesium acetate
100 μg/ml BSA

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μg bacteriophage λ (Hind III digestion) at 25°C for 1 hr in 50-μl reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® Buffer IV	1 X	5 μl
Apa I	50 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 25°C for 1 hr

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® FastCut Buffer	1 X	5 μl
Apa I	50 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 25°C for 15 min

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.