



Restriction Enzyme BamH I



Cat.#	Size	Conc.
FG-BamHI	10,000 units	20 units/μl

Store at -20°C

Supplied with: 10X FastGene® Buffer BamH I (FG-REBBamHI)
10X FastGene® FastCut Buffer (FG-REBHF)
6X DNA Loading Buffer
Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Heat Inactivation

No.

Methylation sensitivity

dam methylation: Not sensitive
dcm methylation: Not sensitive
CpG methylation: Not sensitive

Prolonged incubation

A minimum amount of enzyme required to digest 1 μg substrate DNA for 16 hr; 0.5 U.

Relative activity in FastGene® Buffers

FastGene® Buffer I:	75%
FastGene® Buffer II:	100%
FastGene® Buffer III:	100%
FastGene® Buffer IV:	100%
FastGene® FastCut Buffer:	100%

Note

Long-term storage could reduce its catalytic efficiency and specificity. It is not sensitive to *dam*, *dcm*, or mammalian CpG methylation. Reaction condition of low salt, excess enzyme, excess glycerol (>5%) or high pH (>8.0) could result in star activity. Star activity is often observed in NaCl concentrations below 100 mM. It is markedly affected by impurities in DNA. It can cleave DNA with 2 bases on each side of the recognition site.

Source: *Bacillus amyloliquefaciens* H

Reaction conditions

1X FastGene® Buffer BamH I, 37°C
2X FastGene® FastCut Buffer, 37°C

FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 min with FastGene® FastCut Buffer.

1X FastGene® Buffer BamH I

10 mM Tris-HCl (pH 7.9 at 25°C)
150 mM NaCl
10 mM MgCl₂
100 μg/ml BSA

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μg bacteriophage λ at 37°C for 1 hr in 50 μl reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Dilution buffer

FastGene® Diluent A

Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® Buffer BamH I	1 X	5 μl
BamH I	20 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 37°C for 1 hr

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® FastCut Buffer	2 X	10 μl
BamH I	20 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 37°C for 15 min

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.