

Restriction Enzyme BsmB I



Cat.# FG-BsmBI Size 200 units Conc. 10 units/µl

(FG-REB3)

Store at -20℃

Supplied with: 10X FastGene® Buffer III

10X FastGene® FastCut Buffer (FG-REBHF) 6X DNA Loading Buffer

Sterile water

Recognition site

5' ··· C G T C T C (N) ··· 3'

For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Dilution buffer:

FastGene® Diluent B

Heat Inactivation

BsmB I can be inactivated at 80°C for 20 min.

Methylation sensitivity

dam methylation: Not sensitive dcm methylation: Not sensitive CpG methylation: sensitive

Prolonged incubation

A minimum amount of enzyme required to digest 1 μg substrate DNA for 16 hr; 0.5 U.

Relative activity in FastGene® Buffers

FastGene® Buffer I: 10%
FastGene® Buffer II: 50%
FastGene® Buffer III: 100%
FastGene® Buffer IV: 25%
FastGene® FastCut Buffer: 100%

Note

It is an isoschizomer of Esp3 I. Cleavage of mammalian genomic DNA is blocked by CpG methylation.

Source: Bacillus stearothermophilus B61

Reaction conditions

1X FastGene® Buffer III 55℃ 1X FastGene® FastCut Buffer, 55℃

FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

1X FastGene® Buffer III

50 mM Tris-HCl (pH 7.9 at 25°C) 100 mM NaCl 10 mM MgCl $_2$ 100 μ g/ml BSA

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μ g bacteriophage λ at 55°C for 1 hr in 50 μ l reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assayExtreme pure assay

Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	Χ μΙ
10X FastGene® Buffer III	1 X	5 μΙ
BsmB I	10 unit	1 μΙ
Sterile water		up to 50 μl

- → Incubate at 55°C for 1 hr
- Fast protocol

rast protocor		
Component	Final Conc.	Volume
Substrate DNA	1 μg	Χ μΙ
10X FastGene® FastCut Buffer	1 X	5 μΙ
BsmB I	10 unit	1 μΙ
Sterile water		up to 50 μl

→ Incubate at 55°C for 15 min

 $\ensuremath{\mathbb{X}}$ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.