



Restriction Enzyme HinP1 I




Cat.#	Size	Conc.
FG-HinP1I	2,000 units	10 units/µl

Store at -20°C

Supplied with: 10X FastGene® Buffer II (FG-REB2)
10X FastGene® FastCut Buffer (FG-REBHF)
6X DNA Loading Buffer
Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Dilution buffer:

FastGene® Diluent A

Heat Inactivation

HinP1 I can be inactivated at 65°C for 20 min.

Methylation sensitivity

dam methylation: Not sensitive

dcm methylation: Not sensitive

CpG methylation: sensitive

Prolonged incubation

A minimum amount of enzyme required to digest 1 µg substrate DNA for 16 hr; 0.25 U.

Relative activity in FastGene® Buffers

FastGene® Buffer I:	50%
FastGene® Buffer II:	100%
FastGene® Buffer III:	100%
FastGene® Buffer IV:	75%
FastGene® FastCut Buffer:	100%

Note

It is an isoschizomer of Hha I. It produces a 5' extension, whereas Hha I produces a 3' extension. Thus, ligation is more efficient with HinP1 I cleaved fragments. Cleavage of mammalian genomic DNA is blocked by CpG methylation.

Source: *Haemophilus influenzae* P1

Reaction conditions

1X FastGene® Buffer II 37°C

1X FastGene® FastCut Buffer, 37°C

FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

1X FastGene® Buffer II

10 mM Tris-HCl (pH 7.9 at 25°C)

50 mM NaCl

10 mM MgCl₂

100 µg/ml BSA

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 µg bacteriophage λ at 37°C for 1 hr in 50 µl reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene® Buffer II	1 X	5 µl
HinP1 I	10 unit	1 µl
Sterile water		up to 50 µl

→ Incubate at 37°C for 1 hr

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene® FastCut Buffer	1 X	5 µl
HinP1 I	10 unit	1 µl
Sterile water		up to 50 µl

→ Incubate at 37°C for 15 min

※ We recommend 5-10 units of enzyme per µg DNA and 10-20 units for genomic DNA in a 1 h digest.