

# Restriction Enzyme Pvu II



Cat.# Size Conc. FG-Pvull 5,000 units 10 units/μl

Store at -20°C

**Supplied with:** 10X FastGene® Buffer II (FG-REB2)

10X FastGene® FastCut Buffer (FG-REBHF)

6X DNA Loading Buffer

Sterile water

# Recognition site

For Research Use Only. Not for use in diagnostic procedures.

**ISO**9001

### Dilution buffer

FastGene® Diluent B

## **Heat Inactivation**

No.

### Methylation sensitivity

*dam* methylation: Not sensitive *dcm* methylation: Not sensitive CpG methylation: Not sensitive

## Prolonged incubation

A minimum amount of enzyme required to digest 1- $\mu$ g substrate DNA for 16 hr; 0.13 U.

# Relative activity in FastGene® Buffers

 FastGene® Buffer I:
 75%

 FastGene® Buffer II:
 100%

 FastGene® Buffer III:
 25%

 FastGene® Buffer IV:
 10%

 FastGene® FastCut Buffer:
 100%

#### Note

It is not sensitive to *dam, dcm,* or mammalian CpG methylation. Reaction condition of low salt, excess enzyme, excess glycerol (>5%) or high pH (>8.0) may result in star activity.

Source: Proteus vulgaris

### Reaction conditions

1X FastGene® Buffer II, 37°C 1X FastGene® FastCut Buffer, 37°C

### FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 min with FastGene® FastCut Buffer.

### 1X FastGene® Buffer II

10 mM Tris-HCl (pH 7.9 at 25°C) 50 mM NaCl 10 mM MgCl₂ 100 µg/ml BSA

### Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1  $\mu$ g bacteriophage  $\lambda$  at 37°C for 1 hr in 50  $\mu$ l reaction mixtures.

# **Quality control**

- Unit definition assay
- Overdigestion assay
- Endonuclease assay - Extreme pure assay

# Standard reaction condition

- Normal protocol

- Normal protocol		
Component	Final Conc.	Volume
Substrate DNA	1 μg	Χ μΙ
10X FastGene® Buffer II	1 X	5 μΙ
Pvu II	10 unit	1 μΙ
Sterile water		up to 50 μl
→ Incubate at 37°C for 1 hr		

Component	Final Conc.	Volume
Substrate DNA	1 μg	Χ μΙ
10X FastGene® FastCut Buffer	1 X	5 μΙ
Pvu II	10 unit	1 μΙ
Sterile water		up to 50 μl

→ Incubate at 37°C for 15 min

 $\ensuremath{\mathbb{X}}$  We recommend 5-10 units of enzyme per  $\mu g$  DNA and 10-20 units for genomic DNA in a 1 h digest.