FastGene®

Restriction Enzyme Sbf I

IV 37° 80°

Cat.#	Size	Conc.
FG-SbfI	500 units	10 units/µl

Store at -20℃

Supplied with: 10X FastGene® Buffer IV (FG-REB4)
10X FastGene® FastCut Buffer (FG-REBHF)
6X DNA Loading Buffer
Sterile water

Recognition site

For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Source: *Streptomyces* species Bf-61

Reaction conditions
1X FastGene® Buffer IV, 37℃
1X FastGene® FastCut Buffer, 37℃

FastGene® FastCut Buffer
FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

1X FastGene® Buffer IV
20 mM Tris-acetate (pH 7.9 at 25℃)
50 mM potassium acetate
10 mM magnesium acetate
100 µg/ml BSA

Unit definition
One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37℃ in a total reaction volume of 50 µl.

Quality control
- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Dilution buffer:
FastGene® Diluent A

Heat Inactivation
Sbf I can be inactivated at 80℃ for 20 min.

Methylation sensitivity
dam methylation: Not sensitive
dcm methylation: Not sensitive
CpG methylation: Not sensitive

Relative activity in FastGene® Buffers

FastGene® Buffer I:	50%
FastGene® Buffer II:	25%
FastGene® Buffer III:	10%
FastGene® Buffer IV:	100%
FastGene® FastCut Buffer:	100%

Note
It is an isoschizomer of Sse8387 I. Reaction condition with excess enzyme, excess glycerol (>5%) or high pH (>8.0) may result in star activity.

Standard reaction condition
- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene® Buffer IV	1 X	5 µl
Sbf I	10 unit	1 µl
Sterile water		up to 50 µl

→ Incubate at 37℃ for 1 hr

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene® FastCut Buffer	1 X	5 µl
Sbf I	10 unit	1 µl
Sterile water		up to 50 µl

→ Incubate at 37℃ for 15 min

※ We recommend 5-10 units of enzyme per µg DNA and 10-20 units for genomic DNA in a 1 h digest.