

# Acetamide Broth Base ISO

Cat. 1155

For confirmation of *Pseudomonas aeruginosa* by membrane filtration

## Practical information

Industry: Water

Regulations: ISO 16266

## Principles and uses

Acetamide Broth Base ISO contains acetamide which is the sole source of carbon. It is used for the confirmation and identification of *Pseudomonas aeruginosa*, as specified by the ISO 16266. It uses the ability of non-fermenting Gram-negative bacteria to deaminate the acetamide. The deamination of the acetamide produces ammonia which increases the pH of the medium, acetamide deamination is accomplished by *P. aeruginosa*, *P. acidovorans*, Group III (*Achromobacter xylosoxidans*), and *Alcaligenes odorans*.

Acetamide is the single carbon source. The Potassium salt has a high buffering capacity and Sodium chloride supplies essential electrolytes for transport and osmotic balance.

It is prepared according to ISO 16266.

*Pseudomonas aeruginosa* is an opportunist pathogen for humans, capable of growing in water with a low concentration of nutrients. This is why natural mineral water and spring water are *Pseudomonas aeruginosa* free at the time of their commercialization. This microorganism can also be found in swimming pool water.

## Formula in g/L

Acetamide	2	Magnesium sulfate	0,2
Monopotassium phosphate	1	Sodium chloride	0,2

## Preparation

Suspend 3,4 grams of the medium in 900 ml of distilled water. Adjust the pH to  $7,0 \pm 0,5$  at 25 °C. Add one ml of recently prepared Solution B. Whilst agitating add water to obtain a final volume of one liter. Distribute into tubes in 5 ml aliquots, close and sterilize in autoclave at 121 °C for 15 minutes. Prepared tubes must be stored in a dark place.

## Instructions for use

According to ISO 16266 for the detection and enumeration of *Pseudomonas aeruginosa*:

- Filter a certain volume of water sample through a filter membrane and place the membrane on a *Pseudomonas* CN Agar Base plate (Cat. 1153).
- Incubate at a temperature of  $36 \pm 2$  °C for  $44 \pm 4$  h.
- Count the colonies that have a green/blue pigmentation (pyocyanin) as confirmed *P. aeruginosa*.
- Examine the membrane under UV light.
- All colonies that are fluorescence (+) and reddish-brown colonies should be confirmed.
- Spread all the colonies that should be confirmed on Nutrient Agar plates (Cat. 1156) to obtain pure cultures. Incubate at  $36 \pm 2$  °C for  $22 \pm 2$  h.
- Perform oxidase assay to the reddish-brown colonies.
- Streak the oxidase (+) colonies on King B Medium (Cat. 1154) to check the fluorescence production. Incubate at  $36 \pm 2$  °C for up to 5 days. Normally 24 hours are enough.
- Inoculate all the fluorescence (+) colonies, both in CN agar and in King B Medium, in the Acetamide Broth (Cat. 1155 or Cat. 2017) medium and add one or two drops of Nessler reagent to check the ammonia production. Incubate at  $36 \pm 2$  °C for  $22 \pm 2$  h.
- The colonies that produce pyocyanin in CN agar, the colonies fluorescence (+) in CN agar and ammonia (+) in Acetamide broth, and the reddish brown colonies in CN agar, oxidase (+), fluorescence (+) in King B Agar and ammonia (+) in Acetamide Broth, are counted as confirmed *P. aeruginosa*.

## Quality control

Solubility	Appearance	Color of the dehydrated medium	Color of the prepared medium	Final pH (25°C)
------------	------------	--------------------------------	------------------------------	-----------------

## Microbiological test

---

Incubation conditions: (36±2 °C / 22±2 h)

Microrganisms	Specification	Characteristic reaction
Pseudomonas aeruginosa ATCC 10145	Good growth	Ammonium production
Pseudomonas aeruginosa ATCC 27853	Good growth	Ammonium production
Pseudomonas aeruginosa ATCC 9027	Good growth	Ammonium production

## Storage

---

Temp. Min.:2 °C  
Temp. Max.:25 °C

## Bibliography

---

ISO 16266 Water quality -- Detection and enumeration of Pseudomonas aeruginosa -- Method by membrane filtration