# visocolor® ECO Phosphate

en

Test kit for performing colorimetric tests on phosphate ions in surface water and sewage

#### Method:

Ammonium molybdate forms with phosphate ions phosphomolybdic acid, which is reduced to phosphomolybdenum blue.

#### Measurement range:

0.2-5 mg/L PO<sub>4</sub>-P

## Contents of test kit (\*refill pack):

sufficient for 80 tests

25 mL PO<sub>4</sub>-1\* 25 mL PO<sub>4</sub>-2\*

2 screw-plug measuring glasses

1 slide comparator

1 colour chart

1 plastic syringe 5 mL

1 instructions for use'

#### **Hazard warning:**

Information regarding safety can be found on the box' label and in the safety data sheet. You can download the SDS from www.mn-net.com/SDS.

#### Instructions for use:

## a) colorimetric determination with color chart

also refer to the pictogram on the back of the color chart

 Pour a 5 mL water sample into each of the measuring glasses using the plastic syringe.
Place a measuring glass on position A in the comparator.

## Only add the reagent to measuring glass B.

- 2. Add 6 drops of PO<sub>4</sub>-1, seal the glass and mix.
- 3. Add 6 drops of PO<sub>4</sub>-2, seal the glass and mix.
- 4. Open the glass after 10 min and place it on position B in the comparator.
- Slide the comparator until the colors match in the inspection hole on top. Check the measurement reading in the recess on the comparator reed. Mid-values can be estimated.
- 6. After use, rinse out both measuring glasses thoroughly and seal them.

## b) photometric determination

The reagents are also suitable for **photometric evaluation.** Please refer to the separate instructions for photometric performance.

This technique can be used also for analyzing sea water.

# Disposing of the samples:

Information regarding disposal can be found in the safety data sheet. You can download the SDS from www.mn-net.com/SDS.

## Interferences:

Larger amounts of oxidizing reagents inhibit formation of the blue color complex and have to be destroyed.  $\rm H_2S$  interferes in concentrations above 2 mg/L, but can be expelled after acidification of the water sample. Heavy metals in excess of 10 mg/L can slightly decrease the intensity of the color (vanadium causes an increase in color). Silica interferes in excess of 10 mg/L Si.

# Conversion table:

mg/L PO <sub>4</sub> -P	mg/L PO <sub>4</sub> 3-	$mg/L P_2O_5$
0.2	0.6	0.5
0.3	0.9	0.7
0.5	1.5	1.1
0.7	2.1	1.6
1	3	2
2	6	5
3	9	7
5	15	12

## Storage

Store the test kit in a cool (< 25 °C) and dry place.