

Dulbecco's Phosphate Buffered Saline w/o Calcium w/o Magnesium

CAT N° : P0750

Theoretical Ph : 7.5 ± 0.3

Osmolality : $285 \text{ mOsm/kg} \pm 10\%$

Colour : Off-white powder

Storage conditions : Room temperature

Shelf life : 48 months

Endotoxin : $< 1 \text{ EU/ml}$

Composition : Displayed on website; also available on request

Recommended use :

- Respect storage conditions of the product
- Do not use the product after its expiry date
- Store the product in a dry area
- Wear clothes adapted to the manipulation of the product to avoid contamination (e.g. : gloves, mask, hygiene cap, overall...)
- Protect the product from any form of humidity
- Use, in one time, after opening, the entire quantity of product of the container, without making a concentrated solution (to avoid the formation of precipitates). If it is not possible, close the container immediately after sampling the quantity of powder required.
- Supplements can be added prior to sterile filtration of the medium or aseptically introduced to sterile medium (respect the final concentration of the media). The nature of the supplements may affect storage conditions and shelf life of the medium.

The product is intended to be used *in vitro*, in laboratory only. Do not use it in therapy, human or veterinary applications.

Application :

Dulbecco's Phosphate Buffered Saline (DPBS) is intended for use in the maintenance of mammalian cells where a chemically defined, balanced salt solution provides an environment that will maintain the structural and physiological integrity of cells *in vitro*. DPBS is commonly used in cell enumerations as a diluent, for rinsing cells and as a buffer in many chromatographic procedures. DPBS is also used to wash and resuspend cells during the dissociation process. This product has neither calcium nor magnesium nor sodium bicarbonate ions in its composition so it supports the activity during trypsinisation.

Preparation instructions :

- 1) Measure out 90% of final required volume of water. Water temperature should be $15\text{-}20^\circ\text{C}$.
- 2) While gently stirring the water, add the powdered medium (9.55g/l). Stir until dissolved. Do not heat.
- 3) Rinse original package with a small amount of water to remove all traces of powder. Add to solution in step 2.
- 4) While stirring, adjust the pH of the medium to 0.1-0.3 pH units below the desired pH since it may rise during filtration. The use of 1 N HCl or 1 N NaOH is recommended.
- 5) Add additional water to bring the solution to final volume.
- 6) Sterilize immediately by filtration using a membrane with a porosity of 0.22 microns.
- 7) Aseptically dispense medium into sterile container.

Indications of deterioration :

Dry powder medium should be free flowing. Do not use if powder caked. Prepared medium should be cleared of particulates and flocculent material. Do not use if liquid medium is cloudy or contains precipitate. Other evidence of deterioration may include color change or degradation of physical or performance characteristics.

DOMINIQUE DUTSCHER SAS