ExoProStar S

ENZYMATIC PCR AND SEQUENCE REACTION CLEANUP

ExoProStar™ S is optimized to purify PCR and sequencing setup reactions quickly, efficiently, and reliably.

ExoProStar S contains Amersham™ Shrimp Alkaline phosphatase and Exonuclease I, formulated to work together to remove unincorporated primers and nucleotides from amplification reactions in preparation for sequencing, cloning, genotyping or further DNA modification reactions.

- Enzymes optimized to work together for high efficiency removal of unincorporated primers and nucleotides
- Enzymes provided in two separate tubes, just two simple pipetting steps are needed to prepare the reaction
- Fast 15 min protocol
- Scalable for different reaction sizes
- No loss of PCR product
- Easy to automate
- Complete heat inactivation of the enzymes within 15 min




Fig 1. Schematic representation of the PCR cleanup process using ExoProStar S.

DOMINIQUE DUTSCHER SAS

Optimized for efficient primer digestion

ExoProStar S has been optimized for highly efficient primer digestion, helping to improve the quality of downstream analysis.

No loss of PCR product

The use of an enzymatic digestion approach to clean up amplification reactions reduces losses of PCR product. The process has no intermediate transfer steps, spin columns or binding matrix to retain your PCR product, and double-stranded DNA is left intact by the Exonuclease I and Alkaline Phosphatase enzymes. The size of the PCR fragment does not affect the cleanup efficiency of the reaction (Fig 2).




Fig 2. Agarose gel electrophoresis of different size PCR products pre- and post-digestion with ExoProStar S. The samples were digested for 5 min at 37°C followed by denaturation at 80°C for 10 min according to kit protocol. We did not detect any loss of PCR product in any of the samples.

High-quality sequencing results

Removal of unincorporated primers and nucleotides is essential to high-quality DNA sequencing. Failure to fully remove these components leads to high background signals and miscalling of bases. With ExoProStar S, Phred20 quality scores were routinely achieved at read lengths > 800 bp, equivalent to or better than other approaches to sample preparation (Fig 3).




Fig 3. The importance of sample clean up before DNA sequencing is illustrated in the comparison between panel (A), showing PCR sequence quality following treatment with ExoProStar S and panel (B) showing sequence quality without this treatment. Read length, base calling and sequence quality are significantly improved by the use of ExoProStar S.

Heat inactivation of ExoProStar S enzymes

Downstream operations can be adversely affected by the presence of active Exonuclease I or Alkaline Phosphatase in the PCR product following digestion. It is therefore essential that the enzymes are effectively denatured during the post-digestion heating step.

Some alkaline phosphatase enzymes are tolerant of high temperature treatment and may retain some activity causing problems in later processes. Shrimp Alkaline Phosphatase has been optimized to be quickly denatured, reducing the risk of downstream interference (Fig 4).




Fig 4. Temperature denaturation profile of Shrimp Alkaline Phosphatase at 65°C showing rapid and complete denaturation within 10 min. ExoProStar S protocol recommends denaturation of the enzyme components at 80°C, providing greater confidence in the inactivation of both enzymes prior to further downstream processes.

Kit components and storage

ExoProStar S kit contains one tube of Exonuclease I and one tube of Shrimp Alkaline Phosphatase. The kit is supplied on dry ice and should be stored at -20°C. Enzymes can be sub-aliquoted if required for storage convenience and should be maintained on ice during reaction setup.

DOMINIQUE DUTSCHER SAS

Ordering information

ExoProStar S Enzymatic PCR and Sequence Reaction Cleanup Kit

Quantity	Code number
20 reactions	US79002
100 reactions	US79010
500 reactions	US79050
2000 reactions	US79200
5000 reactions	US79500

Related products

Amplification

Product	Quantity	Code number
dNTP set (100 mM each A,C,G,T)	4 × 100 µmol	28-4065-52
Amersham Ready-To-Go™ RT-PCR Beads (0.2 ml hinged tube with cap)	96 reactions	27-9259-01
PuReTaq Ready-To-Go PCR Beads (0.2 ml hinged tube with cap)	96 reactions	27-9559-01
Amersham Hot Start Mix RTG™ (0.2 ml tubes, 12 × 8 strip wells)	96 reactions	28-9006-53
Taq DNA Polymerase (cloned)	4 × 250 units	27-0798-05

DNA labeling

Cy™5 dUTP	250 nmol	PA55032
Cy3 dUTP	250 nmol	PA53032
Cy5 dCTP	250 nmol	PA55031
Cy3 dCTP	250 nmol	PA53031
CyDye™ Post-Labeling Reactive Dye Pack	12 × Cy3 12 × Cy5	RPN5661

DNA purification

Blood genomicPrep Mini Spin Kit	50	28-9042-64
Tissue and cells genomicPrep Mini Spin Kit	50	28-9042-75
Bacteria genomicPrep Mini Spin Kit	50	28-9042-58

DNA cleanup

GFX™ PCR DNA and Gel Band Purification Kit	100 purifications	28-9034-70
GFX 96 PCR Purification Kit	10 × 96 well plates	28-9034-45
Amersham MicroSpin™ S-400 HR columns	50	27-5140-01
Amersham MicroSpin S-300 HR columns	50	27-5130-01

Enzymes

Amersham Shrimp Alkaline Phosphatase	500 units	E70092Y
Exonuclease I	2500 units	E70073Z

DOMINIQUE DUTSCHER SAS

cytiva.com/ExoProStarS

Cytiva and the Drop logo are trademarks of Global Life Sciences IP Holdco LLC or an affiliate. Amersham, Cy, CyDye, ExoProStar, GFX, MicroSpin, Ready-To-Go, RTG, and Typhoon are trademarks of Global Life Sciences Solutions USA LLC or an affiliate doing business as Cytiva.

The purchase of CyDye products includes a limited license to use the CyDye products for internal research and development but not for any commercial purposes. Cy and CyDye are trademarks of Global Life Sciences Solutions USA LLC or an affiliate doing business as Cytiva. A license to use the Cy and CyDye trademarks for commercial purposes is subject to a separate license agreement with Cytiva. Commercial use shall include:

1. Sale, lease, license or other transfer of the material or any material derived or produced from it.
2. Sale, lease, license or other grant of rights to use this material or any material derived or produced from it.
3. Use of this material to perform services for a fee for third parties, including contract research and drug screening.

If you require a commercial license to use the Cy and CyDye trademarks, please contact LSIicensing@cytiva.com.

© 2020 Cytiva

All goods and services are sold subject to the terms and conditions of sale of the supplying company operating within the Cytiva business. A copy of those terms and conditions is available on request. Contact your local Cytiva representative for the most current information.

For local office contact information, visit cytiva.com/contact

CY14464-16Sep20-DF

cytiva